


Acta Universitatis Szegediensis

Visit us at  
[www.sci.u-szeged.hu/ABS](http://www.sci.u-szeged.hu/ABS)

# Acta Biologica Szegediensis

Volume 49, Number 1-2, 2005



University of Szeged, Szeged, Hungary

## Acta Biologica Szegediensis

Acta Biologica Szegediensis (ISSN 1588-385X print form; ISSN 1588-4082 online form), a member of the Acta Universitatis Szegediensis family of scientific journals (ISSN 0563-0592), is published yearly by the University of Szeged. Acta Biologica Szegediensis covers the growth areas of modern biology and publishes original research articles and reviews, involving, but not restricted to, the fields of anatomy, embryology and histology, anthropology, biochemistry, biophysics, biotechnology, botany and plant physiology, all areas of clinical sciences, conservation biology, ecology, genetics, microbiology, molecular biology, neurosciences, paleontology, pharmacology, physiology and pathophysiology, and zoology. Occasionally, Acta Biologica Szegediensis will publish symposium materials. Acta Biologica Szegediensis particularly encourages young investigators and clinicians to submit novel results of interest.

---

Editor-in-Chief: Károly Gulya

Senior Editors: Dénes Budai (*Cell Physiology*)  
Julius Gy. Papp (*Pharmacology*)  
István Raskó (*Genetics*)

Editorial Board:	Magdolna Ábrahám-Gulyás ( <i>Biochemistry</i> )	Péter Maróti ( <i>Biophysics</i> )
	Mihály Boros ( <i>Experimental Surgery</i> )	Péter Maróy ( <i>Genetics</i> )
	László Erdei ( <i>Plant Physiology</i> )	Erzsébet Mihalik ( <i>Botany</i> )
	Gyula Farkas ( <i>Anthropology</i> )	András Mihály ( <i>Anatomy, Embryology, Histology</i> )
	László Gallé ( <i>Ecology</i> )	Attila Pál ( <i>Obstetrics and Gynecology</i> )
	Zoltán Janka ( <i>Psychiatry</i> )	Aurél J. Simonka ( <i>Traumatology, Surgery</i> )
	Csaba Vágvölgyi ( <i>Microbiology</i> )	József Toldi ( <i>Comparative Physiology</i> )
	Kornél Kovács ( <i>Biotechnology</i> )	László Vécsei ( <i>Neurology</i> )
	János Lonovics ( <i>Internal Medicine</i> )	László Vígh ( <i>Biochemistry</i> )

Technical Editors: Lidia Bakota, Tamás Mikola

---

### Submission of manuscripts

Manuscripts should be prepared in accordance with the Instructions to Authors published in each issue, also available at <http://www.sci.u-szeged.hu/ABS>, and submitted to:

Correspondence relating to the status of the manuscripts, proofs, publication, reprints and advertising should be sent to:

Károly Gulya  
Acta Biologica Szegediensis, Editorial Office  
Department of Zoology and Cell Biology  
University of Szeged  
2 Egyetem u., POB 659, H-6722 Szeged, Hungary  
Phone: 36 (62) 544-048, fax: 36 (62) 544-049  
E-mail: [gulyak@bio.u-szeged.hu](mailto:gulyak@bio.u-szeged.hu)

Lidia Bakota  
Acta Biologica Szegediensis, Editorial Office  
Department of Zoology and Cell Biology  
University of Szeged  
2 Egyetem u., POB 659, H-6722 Szeged, Hungary  
Phone: 36 (62) 544-125, fax: 36 (62) 544-049  
E-mail: [bakotal@bio.u-szeged.hu](mailto:bakotal@bio.u-szeged.hu)

---



### Subscriptions


Acta Biologica Szegediensis is published yearly in four issues per volume. All subscriptions relate to the calendar year and must be pre-paid. The annual subscription rate is currently 50 USD and includes air mail delivery and handling.

Acta Biologica Szegediensis is indexed in BIOSIS Database, EMBASE, Excerpta Medica, Elsevier BIOBASE (Current Awareness in Biological Sciences) and Zoological Record.

The Table of Contents for the current issue and those for previous issues can be found at <http://www.sci.u-szeged.hu/ABS>.

**8<sup>th</sup> Hungarian Congress on Plant Physiology**  
**6<sup>th</sup> Hungarian Conference on Photosynthesis**

**22-25 August, 2005**  
**Szeged, Hungary**



***SUPPORTED BY***

**Hungarian Society for Plant Physiology, member of the FESPB**  
**Photosynthesis "Life from the Light" Foundation**  
**Committee for Plant Physiology, Hungarian Academy of Sciences**  
**Committee for Plant Ecophysiology, Hungarian Academy of Sciences**  
**Biological Research Center, Hungarian Academy of Sciences, Szeged**  
**Department of Plant Physiology, University of Szeged**

**PROCEEDINGS**



**Guest Editor: László Erdei**

## Aspects of *in situ*, *in vitro* germination and mycorrhizal partners of *Liparis loeselii*

Zoltán Illyés\*, Szabolcs Rudnóy, Zoltán Bratek

Department of Plant Physiology and Molecular Plant Biology, Eötvös Loránd University, Budapest, Hungary

**ABSTRACT** Our present work investigated asymbiotic *in vitro* and symbiotic *in situ* germination of the rare and protected terrestrial orchid, *Liparis loeselii* (L.) Rich. as one of the possibilities for its conservation. Asymbiotic germination array was tested on four different asymbiotic media (different macroelement and organic nitrogen levels) in dark and light. Additionally the effect of cold-treatment was also investigated. *In situ* germination rate was moderate, protocorms was observed only in the close surroundings of adult *L. loeselii* individuals. The fungal partners of *L. loeselii* were identified by sequence analysis of the internal transcribed spacer (ITS) of nuclear ribosomal DNA (nrDNA). The 'root segment technique' was effective for isolation of only one species, *Tulasnella* (*Epulorhiza*), whereas using 'in situ baiting' yielded isolates of both *Tulasnella* and *Ceratobasidium* (*Rhizoctonia*) species.

Acta Biol Szeged 49(1-2):137-139 (2005)

### KEY WORDS

*Liparis loeselii*  
germination  
mycorrhiza  
ITS sequence

In most European countries, the populations of fen orchid *Liparis loeselii* L. Rich have dramatically decreased in the past few decades, whereas special efforts of many conservation project have been made for its maintenance and propagation (Ramsay 1994; Vejsadová 2001). Although asymbiotic growing of the species has been studied yet and seedlings have usually been transferred successfully to *ex vitro* conditions, the asymbiotically grown seedlings rarely survive the re-establishment into natural habitats (Takács and Bratek 1999). To gain a deeper insight into autecology and habitat requirements of fen orchid, it is crucial to identify mycorrhizal fungal partners in order to increase the planting survival of symbiotically grown seedlings. Nevertheless, asymbiotic germination can provide both information on ecological requirements of the species and artificial multiplication by producing numerous seedlings. The Hungarian occurrences of *L. loeselii* are unique in Europe, since they live on floating mats (Balogh et al. 1980), in contrary to the rest of Europe, where the species lives on plain fens and temporary marshes. Botanical and mycological investigations may also enhance our knowledge on these unique habitats. DNA-based methods can greatly facilitate the identification of the mycorrhizal fungi, since morphological identification of the fungus partner is generally much more complicated.

### Materials and Methods

#### Asymbiotic germination

The seeds were collected from Lake Velence, where the highest Hungarian population of *L. loeselii* is located (Balogh et al. 2002). Orchid seeds were surface sterilized for 10 min in chloride of lime, washed twice in sterile water and sown on

agar in Petri dishes. The germination was tested on four asymbiotic medium: 'MS' (Murashige and Skoog 1962), 'MS1/2' (modified 'MS' containing half amount of macro elements), 'Debergh' (Van Weas and Debergh 1986) and 'Fast' (Fast 1974). Organic nitrogen resources were included in the latter two media. Cultures were maintained at 25°C under diurnal conditions or in darkness. Some dishes of the latter group were exposed to cold (4-6°C) pre-treatment for one month.

#### *In situ* baiting

Seed packets were constructed from 40x60 mm rectangles of heat-sealed nylon mesh and about 300 seeds of *L. loeselii* were placed in each. The pore size of the mesh (85 µm) was chosen to retain the seeds without impeding fungal growth. The mesh was folded once and clipped into plastic glassless slide mounts (Rasmussen and Whigham 1993). Seed baits were positioned vertically in top-peat (approx. depth 5 cm) on the floating mat of Lake Velence. Each packet was deployed in the reed and willow habitat complex at five localities. Three of the five plots contained adult *L. loeselii* plant(s). Seed baits were buried at two occasions: a) in July 2003 and b) in October 2003 and were removed at three times: a) in October 2003 and b) in June and October 2004.

#### Isolation of fungal symbionts of *Liparis loeselii*

In order to determine the characteristics of fungus forming pelotons in seedlings and roots of *L. loeselii*, symbionts were isolated using aseptic techniques and cultured on Potato Dextrose Agar medium (Hadley 1970). Seedlings taken from packets and roots were surface-sterilized by immersion in 0.1% AgNO<sub>3</sub> for 1-3 min. after washing by distilled water they were placed onto the agar. One of the two adult fen orchids was collected from Lake Velence (Hungary), and the

\*Corresponding author. E-mail: zillyes@ludens.elte.hu






Figure 1. *In vitro* germination 4 months after sowing and *in situ* germination.

other was originated from Ceska Lípa (Czech Republic).

### Molecular identification of fungi

DNA analysis was carried out on 12 fungal strains isolated from *in situ* germinated seedlings and 3 strains isolated from adult *L. loeselii* roots. DNA extraction, PCR and sequencing were carried out according to McKendrick et al. (2000).

### Results and Discussion

The highest germinating rate allied with the fastest germination was observed on 'Fast' and 'Debergh' media, both of which contained organic nitrogen sources. Higher germination rate was detected on the 'MS1/2' medium, comparing to 'MS' (Fig. 1), may be related to the ecological requirements of the plants occurring in oligotrophic habitats. Although the cold pre-treated seeds began to germinate later than the others, after 120 days they showed similar germination rate. The effect of light/dark on germination of temperate terrestrial orchid is considered to be controversial (Arditti et al. 1990; Rasmussen and Rasmussen 1991). Nevertheless, we found that *L. loeselii* seeds germinated on light, but exhibited much lower level of germination than in dark (Fig. 1).

*In situ* germination rate was moderate (Fig. 1) presumably due to low abundance of potential symbionts in the constantly humid fen peat. Other facts also support this scenario: germination was observed 1) in separated groups of seeds in the slide mounts and 2) only in the close surroundings of adult *L. loeselii* plants. *In vitro* germination of seeds at room tem-

perature began four-six weeks after the sowing, in contrast to *in situ* germination, which started germination two months later. The delay of germination might be caused by the cold microclimate of floating mat.

McCormick (2004) found that *L. liliifolia* considered to be the closest relative of *L. loeselii* was associated with a single fungal species, while we isolated likely the same and an additional species from our *L. loeselii* samples. Fungal ITS sequences resembling that of *Tulasnella* (anamorph: *Epulorhiza*) reported from *L. liliifolia*, were isolated from roots of two adult *L. loeselii* individuals (Hungary, Czech Republic) and also from 9 *L. loeselii* protocorms (Table 1). The protocorms also hosted 3 species probably the genus *Ceratobasidium* (anamorph: *Rizoctonia*) (Table 1). Considering the symbiont fungus invariability as for the photosynthetic *L. liliifolia* species, McCormick et al. (2004) rejected the widespread idea that the symbiont specificity only occurs in case of non-photosynthetic plants. Our results support the classical views, whereas the results by McCormick et al. (2004) may be attributed to the poverty in potential symbiont fungal species of the *L. liliifolia* habitats.

The infection of asymbiotically cultivated plants by isolated symbiont fungi can give a huge help to survive for the *L. loeselii* individuals when re-establishing.

### Acknowledgments

We would like to thank Duna-Ipoly National Park and Mr. Gábor Sramkó.

### References

- Arditti J, Ernst R, Yam TW, Glabe C (1990) A contribution of orchid mycorrhizal fungi to seed germination: a speculative review. *Lindleyana* 5:249-255.
- Balogh M, Bratek Z, Illyés Z, Zöld-Balogh Á (2002) A *Liparis loeselii* (L.) Rich. tömeges előfordulása a Velencei-tavon. *Kitaibelia* 7:247.
- Balogh M, Patkó Á, Vári L (1980) An interesting *Liparis* presence and its ecological significance on Lake Velence/Hungary. *Ann Univ Sci Budapest* 22-23:49-55.
- Fast (1974) Über eine Methode der kombinierten generativenvegetativen Vermehrung von *Cypripedium calceolus*. *Die Orchidee* 25:125-129
- Hadley G (1970) Non-specificity of symbiotic infection in orchid mycorrhiza. *New Phytol* 69:1015-1023.
- McCormick MK, Whigham DF, O'Neill J (2004) Mycorrhizal diversity in photosynthetic terrestrial orchids. *New Phytol* 163:425-438.
- McKendrick SL, Leake JR, Taylor DL, Read DJ (2000) Symbiotic germination and development of myco-heterotrophic plants in nature: ontogeny of *Corallorhiza trifida* and characterization of its mycorrhizal fungi. *New Phytol* 145:523-537.

Table 1. Isolated symbionts of *L. loeselii*.

Origin	Isolation source	Genus	No. of strains	Accession no.
Czech Republic	Root	Epulorhiza	2	AJ549128
Hungary	Root	Epulorhiza	1	AJ549124
Hungary	Protocorm	Epulorhiza	9	AM040890
Hungary	Protocorm	Rhizoctonia	3	AM040889

- Murashige T, Skoog F (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol Plant* 155:473-497.
- Ramsay M (1994) Fen orchid re-established in UK. *Kew Scientist* 6.
- Rasmussen HN, Rasmussen RN (1991) Climatic and seasonal regulation of seed plant establishment *Dactylorhiza majalis* inferred from symbiotic experiments *in vitro*. *Lindleyana* 5:221-227.
- Rasmussen HN, Whigham D (1993) Seed ecology of dust seeds *in situ*: a new study technique and its application in terrestrial orchids. *Am J Bot* 80:1374-1378.
- Takács AA, Bratek Z (1999) Experiments for micro-propagation of *Liparis*. In Takács AA, ed., *Liparis* project, Progress report 1998-1999. mscr. Budapest, 36.
- Van Weas J, Debergh PC (1986) *In vitro* germination of some Western European orchids. *Physiol Plant* 67:253-261.
- Vejsadová H (2001) Endangered orchid conservation using *in vitro* method. Planta Europa Conference III. June 2001. Available at [http://www.plantaeuropa.org/html/conference\\_2001/conference\\_poster\\_pres.htm](http://www.plantaeuropa.org/html/conference_2001/conference_poster_pres.htm)